


INDICATION: Imodium A-D controls the symptoms of diarrhéa, including Travelers' Diarrhea. DIRECTIONS: Drink plenty of clear fluids to help prevent dehydration, which may accompany diarrhea. DOSAGE: ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER: Take 2 caplets after the first loose bowel movement and 1 caplet after each subsequent loose bowel movement but no more than 4 caplets a day for no more than 2 days. CHILDREN 9-11 YEARS (60-95) high: Take 1 caplet after the first loose bowel movement and 1/2 caplet after each subsequent loose bowel movement but no more than 3 caplets a day for no more than 3 caplets a day for no more than 2 days. Children under 6 for sowel movement and 1/2 caplet after each subsequent loose bowel movement but no more than 2 caplets a day for no more than 2 days. Children under 6 years old. WARNINGS: DO NOT USE IF POUCH IS OPENED. KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN. Do not use for more than two days unless directed by a physician. DO NOT USE IF DIARRHEA IS ACCOMPANIED BY HIGH FEVER (GREATER THAN 101-67), OR IF BLOOD OR MUCUS IS PRESENT IN THE STOOL, OR IF YOU HAVE HAD A RASH OR OTHER ALLERGIC REACTION TO LOPERAMIDE HCI. If you are taking antibiotics or have a history of liver disease, consult a physician before using this product. As with any drug, if you are pregnant or nursing a baby, seek the advice of a health professional assistance of contact a poison control center immediately. ACTIVE INGREDIENT: Loperamide HCI 2 mg per caplet. INACTIVE INGREDIENTS: Dibasic calcium phosphate, magnesium stearate, microcrystalline cellulose, colloidal silicon dioxide, FD&C BIUE 1 and D&C Yellow 10. Store at 15-30-C (59-86-F). Distributed By: MCNEIL Consumer Products Co., Div. of McNEIL-PPC, Inc. '97 5330690

150% OF SIZE



INDICATION: Imodium X-D controls the symptoms of districts, including freelers. Districts on IRECTIONS: Orinic plenty of clear fillids to help prevent behydration, which may accompany districts. DOSME: ADULTS AND CHILDRES including the property of the pr